

JOURNAL OF NUMBER THEORY 4, 523-531 (1972)

Moments of Class Numbers, III

DIETER WOLKE

*Mathematical Institute, University of Marburg, Marburg, West Germany**Communicated by H. Halberstam*

Received October 9, 1970

By means of the large sieve it is proved that

$$\sum_{n \leq x} h^{\alpha}(-n) = \delta(\alpha) x^{(\alpha+2)/2} + O(x^{(\alpha+2)/2-(1/4)+\epsilon})$$

holds for every $\alpha > 0$ and $\epsilon > 0$ ($h(-n)$ is the number of classes of primitive quadratic forms with discriminant $-n$).

Some applications of the large sieve to certain multiplicative functions are announced.

1. INTRODUCTION

In papers [7] and [8] I gave two slightly different methods of estimating the error term in the asymptotic formula for the sum $\sum_{n \leq x} h^k(-n)$ ($k = 1, 2, \dots$). (For further references, see paper [7].) In [8] the error term was proved to be $\ll x^{[(k+2)/2]-(8-\epsilon)/k}$, if $\epsilon > 0$ and $k \geq k_0(\epsilon)$.

It is the aim of the present paper to improve this result and to extend it to powers not necessarily integral.

Write, as usual,

$$L(s, n) = \sum_{m \geq 1, 2 \nmid m} \left(\frac{-n}{m} \right) m^{-s} \quad (1.1)$$

($\sigma = \operatorname{Re} s > 0$, $(-n/m)$ is the Jacobi symbol).

THEOREM 1. *For every real α and $\epsilon > 0$ we have*

$$\sum_{n \leq x} (L(1, n))^{\alpha} = \eta(\alpha) x + O(x^{3/4+\epsilon}).$$

$\eta(\alpha)$ will be defined in Lemma 1. The constants implied by the symbols $O(\cdot)$ and \ll may depend on α and ϵ .

By Gauss' class-number formula and partial summation one easily gets from Theorem 1,

THEOREM 2. For every $\alpha > 0$ and $\epsilon > 0$ we have

$$\sum_{n \leq x} h^\alpha(-n) = \theta(\alpha) x^{(\alpha+2)/2} + O(x^{(\alpha+2)/2 - (1/4) + \epsilon}).$$

It is highly probable that in Theorem 1 the error term is $\ll x^{0.5+\epsilon}$. The difference depends on the fact that in the large-sieve inequality used below we sum over all primitive characters to certain moduli, whereas for the purpose of Theorem 1 we need estimations for real characters only. The inequalities for real characters proved in Elliott [2] and Wolke [7] do not give anything better.

By ideas similar to those of this paper and of Gallagher [4] one can prove the following:

Let $f(n)$ be a multiplicative arithmetical function with the properties

- (i) $0 \leq f(p^a) \leq C_1 a^{C_2}$ (p prime, $a = 1, 2, \dots$),
- (ii) $\sum_{p \leq x} |f(p) - \tau| \ll x(\ln x)^{-\beta}$ for $\tau = \tau(f) > 5/2$ and every $\beta > 0$.

Then, for $\epsilon > 0$, there exists a $\gamma = \gamma(\epsilon, f) > 0$, such that

$$\sum_{k \leq x^{0.5-\epsilon}} \max_{(k,l)=1} \max_{y \leq x} \left| \sum_{\substack{n \leq y \\ n \equiv l \pmod k}} f(n) - \frac{C(f)}{\phi(k)} \right. \\ \left. \times \prod_{p|k} \left(1 + \frac{f(p)}{p} + \frac{f(p^2)}{p^2} + \dots \right)^{-1} y \ln^{\tau-1} y \right| \ll x(\ln x)^{\tau-\gamma} \quad (1.2)$$

holds.

(1.2) implies

$$\sum_{n \leq x} f(n) d(n-a) = c(f, a) x \ln^\tau x (1 + o(1)). \quad (1.3)$$

($d(n)$ is the divisor function, a is a fixed integer $\neq 0$).

In the special case $f(n) = d_k(n)$ ($k = 3, 4, \dots$) this was shown by Linnik [5]. A detailed proof of (1.2) and (1.3) will appear elsewhere.

2. SOME LEMMAS

In the following let α and $\epsilon > 0$ be fixed and x be sufficiently large, $l_1 = \ln x$, $l_2 = \ln l_1$. For $\sigma > 1$ we have

$$(L(s, n))^\alpha = \prod_{p \geq 2} \left(1 - \left(\frac{-n}{p} \right) p^{-s} \right)^{-\alpha} = \sum_{m=1}^{\infty} \left(\frac{-n}{m} \right) f_\alpha(m) m^{-s}, \quad (2.1)$$

where

$$f_{\alpha}(m) \ll (d(m))^{C(\alpha)}. \quad (2.2)$$

By \mathcal{C} we denote the line from $1 + l_1^{-1} - il_1^2$ to $1 + l_1^{-1} + il_1^2$.

LEMMA 1. *We have*

$$\frac{1}{2\pi i} \int_{\mathcal{C}} \Gamma(s-1) L^{\alpha}(s, n) Z^{s-1} ds = \eta(\alpha)x + O(x^{3/4+\epsilon}),$$

where $Z = x^{3/2}$ and $\eta(\alpha) = \sum_{m=1}^{\infty} \phi(m)/m^3 f_{\alpha}(m^2)$ ($\phi(m)$ is Euler's function).

The proof is similar to that of the corresponding lemma in Barban [1].

LEMMA 2. *Let a_n ($M < n \leq M+N$) be any complex numbers. Then the inequality*

$$\sum_{d \leq Q} \mu^2(d) \left| \sum_{\substack{M < n \leq M+N \\ 2 \nmid n}} a_n \left(\frac{-d}{n} \right) \right|^2 \ll (Q^2 + N) \sum_{M < n \leq M+N} |a_n|^2$$

holds.

This follows immediately from the well-known large sieve inequality for character sums (see Gallagher [3]) and the fact that $\chi(n) = (-d/n)$ is a primitive character modulo either d or $4d$, if d is square-free.

For $y \in (x^{\epsilon}, x]$, $d \in (y, 2y]$, $\mu^2(d) = 1$ (in what follows d always denotes squarefree integers) we write

$$w = \exp(y^{\epsilon/2(|\alpha|+1)}), \quad (2.3)$$

and, for $\sigma \geq 1$,

$$\begin{aligned} L^{\alpha}(s, d) &= \prod_{2 < p \leq y} \times \prod_{y < p \leq w} \times \prod_{p > w} \left(1 - \left(\frac{-d}{p} \right) p^{-s} \right)^{-\alpha} \\ &= L_1(s, d) L_2(s, d) L_3(s, d), \text{ say.} \end{aligned} \quad (2.4)$$

LEMMA 3. *For $s \in \mathcal{C}$ or $s = 1$ we have*

$$L_3(s, d) = 1 + O(y^{-2}).$$

We give the proof in the case $s = 1$.

By the Siegel–Walfisz prime number theorem (see Prachar [6]), we get

$$\begin{aligned} \sum_{p>w} \left(\frac{-d}{p}\right) p^{-1} &= \int_w^\infty \frac{dt}{t^2} \sum_{k=1}^d \left(\frac{-d}{k}\right) (\pi(t, d, k) - \pi(w, d, k)) \\ &= \int_w^\infty \frac{dt}{t^2} \sum_{k=1}^d \left(\frac{-d}{k}\right) \left(\frac{\text{Li } t}{\phi(d)} - \frac{\text{Li } w}{\phi(d)} + O\left(\frac{t}{\phi(d) \ln^B t}\right)\right) \\ &\ll \int_w^\infty \frac{dt}{t \ln^B t} \ll y^{-2}, \quad \text{if } B \geq B_0(\epsilon, \alpha). \end{aligned}$$

Hence

$$\log L_3(1, d) = \alpha \sum_{p>w} \left(\frac{-d}{p}\right) p^{-1} + O(y^{-2}) \ll y^{-2},$$

which is equivalent to the statement of the Lemma.

LEMMA 4. For $s \in \mathcal{C}$ or $s = 1$ we have

$$\sum_{d \in (y, 2y]} |1 - L_2(s, d) L_3(s, d)| \ll y^{3/4+2\epsilon}.$$

Proof. Clearly

$$L_2(s, d) \ll \prod_{p \leq w} \left(1 + \frac{1}{p}\right)^{|\alpha|} \ll \exp\left(|\alpha| \sum_{p \leq w} p^{-1}\right) \ll y^\epsilon \quad (2.5)$$

and

$$\log L_2(s, d) = \alpha \sum_{y < p \leq w} \left(\frac{-d}{p}\right) p^{-s} + O(y^{-1}) = \alpha F(s, d) + O(y^{-1}), \text{ say.} \quad (2.6)$$

In the case $|F(s, d)| \leq 1$, we have

$$L_2(s, d) = 1 + O(|F(s, d)| + y^{-1}).$$

Otherwise we use (2.5): $1 - L_2(s, d) \ll y^\epsilon |F(s, d)|$. Hence, in either case,

$$1 - L_2(s, d) \ll y^\epsilon |F(s, d)| + y^{-1}. \quad (2.7)$$

(2.7) and Lemma 3 imply

$$\sum_{d \in (y, 2y]} |1 - L_2(s, d) L_3(s, d)| \ll y^\epsilon \sum_{d \in (y, 2y]} |F(s, d)| + y^{3/4}. \quad (2.8)$$

We split up the interval $(y, w]$ into $\ll y^\epsilon$ intervals $I_\nu = (u_\nu, u_{\nu+1}]$ with $u_{\nu+1} \leq 2u_\nu$. Writing

$$\left(\sum_{p \in I_\nu} \left(\frac{-d}{p}\right) p^{-s}\right)^2 = \sum_{u_\nu^2 < n \leq u_{\nu+1}^2} \left(\frac{-d}{n}\right) a_n n^{-s},$$

we get, with Hölder's inequality and Lemma 2,

$$\begin{aligned} \sum_{d \in (y, 2y]} |F(s, d)| &\ll y^{3/4} \sum_{\nu} \left(\sum_{d \leq 2y} \left| \sum_{u_{\nu}^2 < n \leq u_{\nu+1}^2} \left(\frac{-d}{n} \right) a_n n^{-s} \right|^2 \right)^{1/4} \\ &\ll y^{3/4} \sum_{\nu} ((y^2 + u_{\nu}^2) u_{\nu}^{-2})^{1/4} \ll y^{3/4+\epsilon}. \end{aligned} \quad (2.9)$$

The Lemma now follows from (2.8) and (2.9).

For

$$5/6 \leq \sigma \leq 1 - \frac{\epsilon}{6} \quad (2.10)$$

let \mathcal{C}_{σ} denote the path which connects the four points $1 + l_1^{-1} - il_1^2$, $\sigma - il_1^2$, $\sigma + il_1^2$, and $1 + l_1^{-1} + il_1^2$ and let \mathcal{C}_{σ}' be the line connecting the points $\sigma - il_1^2$ and $\sigma + il_1^2$. By $A_{\sigma} = A_{\sigma}(y)$ we denote the number of $d \in (y, 2y]$ such that

$$|L_1(s, d)| > y^{\epsilon} \quad (2.11)$$

holds for at least one $s \in \mathcal{C}_{\sigma}$.

LEMMA 5. *We have the inequality*

$$A_{\sigma} \ll y^{6(1-\sigma)+\epsilon}.$$

Proof. Put $y' = (\log y)^{1/(1-\sigma)}$. Because of

$$\prod_{p \leq y'} \left(1 - \left(\frac{-d}{p} \right) p^{-s} \right)^{-\alpha} \ll y^{\epsilon/2} \quad \text{for } s \in \mathcal{C}_{\sigma}$$

it is sufficient to estimate the number of $d \in (y, 2y]$ with

$$|G(s, d)| = \left| \sum_{y' < p \leq y} \left(\frac{-d}{p} \right) p^{-s} \right| > \frac{\epsilon}{3} \ln y \quad \text{for at least one } s \in \mathcal{C}_{\sigma}. \quad (2.12)$$

For simplicity, we estimate only A_{σ}' , the number of $d \in (y, 2y]$ such that (2.12) holds for at least one $s \in \mathcal{C}_{\sigma}'$. The contribution of the horizontal parts of \mathcal{C}_{σ} may be treated in the same manner.

We split up the interval $(y', y]$ into $\ll l_1$ intervals $(z_{\nu}, z_{\nu+1}]$ with $z_{\nu+1} \leq z_{\nu}$. The interval $[-l_1^2, l_1^2]$ we dissect into $\ll l_1^3$ parts $[w_{\mu}, w_{\mu+1}]$ with $|w_{\mu} - w_{\mu+1}| \leq l_1^{-1}$. Write

$$T_{\nu}(s, d) = \sum_{z_{\nu} < p \leq z_{\nu+1}} \left(\frac{-d}{p} \right) p^{-s}. \quad (2.13)$$

For $\nu \ll l_1$, $\mu \ll l_1^3$, and $0 \leq \rho \leq l_1^4$ let $A'_{\nu, \mu, \rho} = A'_{\nu, \mu, \rho}(\sigma, y)$ be the number of $d \in (y, 2y]$ for which

$$|T_v^{(\rho)}(\sigma + iw_\mu, d)| \geq (\ln z_{\nu+1})^{\rho-2} \quad (2.14)$$

holds. By Taylor's formula one easily sees that $|G(s, d)| \ll 1$ is true for all $s \in \mathcal{C}_\sigma'$ if d is not counted in any $A'_{\nu\mu\rho}$. Hence, by (2.12),

$$A_\sigma' \leq \sum_{\nu, \mu, \rho} A'_{\nu, \mu, \rho}. \quad (2.15)$$

Let ν, μ and ρ be fixed and write $z = z_{\nu+1}$. Clearly

$$A'_{\nu, \mu, \rho} \leq ((\ln z)^{\rho-2})^{-2r} \sum_{d \leq 2y} |T_v^{(\rho)}(\sigma + iw_\mu, d)|^{2r},$$

where

$$r = [2 \ln y / \ln z] + 1. \quad (2.16)$$

Obviously,

$$(T_v^{(\rho)}(s, d))^r = \sum_{z_\nu^r < n \leq z_{\nu+1}^r} \left(\frac{-d}{n}\right) n^{-s} a_n$$

with

$$a_n = \sum_{\substack{n=p_1 \cdots p_r \\ p_j \in (z_\nu, z_{\nu+1}] \\ j=1, \dots, r}} \ln^\rho p_1 \cdots \ln^\rho p_r \ll C^r r! (\ln z)^{\rho r}$$

and

$$\sum_n a_n \leq \left(\sum_{z_\nu < p \leq z_{\nu+1}} \ln^\rho p \right)^r \ll (\ln z)^{\rho r} z^r.$$

The last lines and Lemma 2 imply

$$A'_{\nu, \mu, \rho} \ll C^r r! (\ln z)^{4r} z^{2r(1-\sigma)}. \quad (2.17)$$

In the case $z \in (y', y^*]$ one easily verifies the inequalities

$$C^r \ll y^{\epsilon/4}, \quad r! \ll y^{2(1-\sigma)+\epsilon/4}, \quad \text{and} \quad z^{2r(1-\sigma)} \ll y^{4(1-\sigma)+\epsilon/4}.$$

In the case $z \in (y^\epsilon, y]$, we have

$$C^r r! \ll 1 \quad \text{and} \quad z^{2r(1-\sigma)} \leq z^{(4(\ln y / \ln z) + 2)(1-\sigma)} \leq y^{6(1-\sigma)}.$$

Hence, in both cases,

$$A'_{\nu, \mu, \rho} \ll y^{6(1-\sigma)+(3/4)\epsilon}. \quad (2.18)$$

Using (2.15), we finally get the required result.

3. PROOF OF THEOREM 1

Every n can be written uniquely in the form $n = dR^2Q^2$ with $\mu^2(d) = 1$, $p \mid R \Rightarrow p \mid d$, and $(Q, d) = 1$. Obviously

$$L^\alpha(s, n) = L^\alpha(s, d) F_Q(s), \quad (3.1)$$

where $F_Q(s)$ is regular and $\ll x^\epsilon$ for $\sigma \geq \frac{1}{2}$. Hence

$$\sum_{n \leq x} L^\alpha(s, n) = \sum_{d \leq x} L^\alpha(s, d) G_d(s), \quad (3.2)$$

where, with the above notation,

$$G_d(s) = \sum_{R^2 Q^2 \leq x/d} F_Q(s) \ll x^{1/2+\epsilon} d^{-1/2}. \quad (3.3)$$

For a fixed integer $\kappa \geq \epsilon^{-1}$ we write

$$\sigma_j = \frac{5}{6} + \frac{j}{6\kappa} \quad (j = 0, 1, \dots, \kappa - 1); \quad (3.4)$$

\sum_ν stands for summation over all intervals

$$J_\nu = (y_{\nu+1}, y_\nu] = (x2^{-\nu-1}, x2^{-\nu}]$$

with $\nu \geq 0$ and $x2^{-\nu} \geq x^\epsilon$.

For every ν we split up the set $\{d \mid d \in J_\nu\}$ into disjoint subsets $\mathcal{S}_{\nu,0}, \dots, \mathcal{S}_{\nu,j-1}$, such that the following is valid:

$$|L_1(s, d)| \leq y_\nu^\epsilon \quad \text{if } d \in \mathcal{S}_{\nu,j}, \quad s \in \mathcal{C}_{\sigma_j}, \quad \text{and } 0 \leq j \leq \kappa - 2. \quad (3.5)$$

By Lemma 5 one sees that the sets $\mathcal{S}_{\nu,j}$ may be chosen in such a way that

$$S_{\nu,j} = \text{card}(\mathcal{S}_{\nu,j}) \ll y_\nu^{6(1-\sigma_j)+2\epsilon} \quad (3.6)$$

holds for $1 \leq j \leq \kappa - 1$. In particular,

$$S_{\nu,\kappa-1} \ll y_\nu^{3\epsilon} \quad \text{for every } \nu. \quad (3.7)$$

We mention Siegel's result (see Prachar [6])

$$L^\alpha(s, n) \ll x^\epsilon, \quad \text{if } n \leq x \text{ and } s \in \mathcal{C} \text{ or } s = 1. \quad (3.8)$$

Because of (3.2) and (3.3) we can write, with $Z = x^{3/2}$,

$$\begin{aligned} I(x) &= \frac{1}{2\pi i} \sum_{n \leq x} \int_{\mathcal{C}} \Gamma(s-1) L^\alpha(s, n) Z^{s-1} ds \\ &= \frac{1}{2\pi i} \sum_{\nu} \int_{\mathcal{C}} \Gamma(s-1) Z^{s-1} \sum_{d \in J_{\nu}} L^\alpha(s, d) G_d(s) ds \\ &\quad + O\left(x^{0.5+2\epsilon} \sum_{d \leq x^\epsilon} \int_{\mathcal{C}} |\Gamma(s-1) L^\alpha(s, d)| |ds|\right), \end{aligned}$$

and further, by (2.4) and (3.8),

$$\begin{aligned} I(x) &= \frac{1}{2\pi i} \sum_{\nu} \int_{\mathcal{C}} \Gamma(s-1) Z^{s-1} \sum_{d \in J_{\nu}} L_1(s, d) G_d(s) ds \\ &\quad + O\left(x^{1/2+\epsilon} \sum_{\nu} y_{\nu}^{-1/2} \int_{\mathcal{C}} |\Gamma(s-1)| \right. \\ &\quad \left. \times \sum_{d \in J_{\nu}} |L_1(s, d)| |1 - L_2 L_3(s, d)| |ds|\right) + O(x^{3/4}). \quad (3.9) \end{aligned}$$

By Lemma 4 and a trivial estimation for $L_1(s, d)$ the first error term is $\ll x^{1/2+4\epsilon} \sum_{\nu} y_{\nu}^{1/4} \ll x^{3/4+4\epsilon}$. For $d \in \mathcal{S}_{\nu, k-1}$ we estimate the integral trivially. Hence, by (3.7) and (3.8), the contribution of all sets $\mathcal{S}_{\nu, k-1}$ is $\ll x^{3/4}$.

For $d \in \mathcal{S}_{\nu, j}$ ($0 \leq j \leq \kappa - 2$) we replace the integral over \mathcal{C} by the integral over \mathcal{C}_{σ_j} . The pole of the Γ -function at the point $s = 0$ produces the residue $L_1(1, d) G_d(1)$. Collecting the last results, we get

$$\begin{aligned} I(x) &= \sum_{\nu} \sum_{0 \leq j \leq \kappa-2} \sum_{d \in \mathcal{S}_{\nu, j}} L_1(1, d) G_d(1) \\ &\quad + O\left(x^{1/2+\epsilon} \sum_{\nu} y_{\nu}^{-1/2} \sum_{0 \leq j \leq \kappa-2} Z^{\sigma_j-1} \int_{\mathcal{C}_{\sigma_j}} \sum_{d \in \mathcal{S}_{\nu, j}} |L_1(s, d)| |ds|\right) \\ &\quad + O(x^{3/4+4\epsilon}). \quad (3.10) \end{aligned}$$

By calculations similar to those above one can show that the main term on the right side of (3.10) is

$$= \sum_{n \leq x} L^\alpha(1, n) + O(x^{3/4+4\epsilon}).$$

To the first error term we apply (3.5), (3.6), and (3.4):

$$\begin{aligned}
 I(x) - \sum_{n \leq x} L^{\alpha}(1, n) &\ll x^{3/4+4\epsilon} + x^{1/2+4\epsilon} \sum_{\nu} y_{\nu}^{-1/2} \sum_{0 \leq j \leq \kappa-2} x^{3/2(\sigma_j-1)} y_{\nu}^{6(1-\sigma_j)} \\
 &\ll x^{3/4+4\epsilon} + x^{1/2+4\epsilon} \sum_{\nu} \sum_{0 \leq j \leq \kappa-2} y_{\nu}^{9/2(1-\sigma_j)-1/2} \\
 &\ll x^{3/4+4\epsilon} + x^{1/2+4\epsilon} \sum_{\nu} y_{\nu}^{1/4} \ll x^{3/4+5\epsilon}.
 \end{aligned}$$

This, in connection with Lemma 1, proves Theorem 1.

REFERENCES

1. M. B. BARBAN, Linnik's 'large sieve' and a limit theorem for the class number of ideals of an imaginary quadratic field, *Izv. Akad. Nauk. SSSR Ser. Mat.* **26** (1962), 573-580.
2. P. D. T. A. ELLIOTT, On the mean value of $f(p)$, *Proc. London Math. Soc.* **21** (1970), 28-96.
3. P. X. GALLAGHER, On the large sieve, *Mathematika* **14** (1967), 14-20.
4. P. X. GALLAGHER, Bombieri's mean value theorem. *Mathematika* **15** (1968), 1-6.
5. JU. V. LINNIK, "The Dispersion Method in Binary Additive Problems," American Mathematical Society, Providence, RI, 1963.
6. K. PRACHAR, "Primzahlverteilung," Springer, Berlin, 1957.
7. D. WOLKE, Moments of the number of classes of primitive quadratic forms with negative discriminant, *J. Number Theory* **1** (1969), 502-511.
8. D. WOLKE, Moments of class numbers, II, to appear in *Archiv der Mathematik*.